

Critical Care

Antifungal Portfolio

CONCORD BIOTECH

Sr. No.	Brand Name	Molecule	SKUs	INDICATION*	DOSE*	PACKSHOT
1	Amfoterol	Liposomal Amphotericin-B	Amfoterol Injection 50 mg	Empirical therapy for presumed fungal infection in febrile, neutropenic patients Cryptococcal Meningitis in HIV infected patients Systemic fungal infections: Aspergillus, Candida, Cryptococcus Treatment of Visceral leishmaniasis	Empirical therapy - 3mg / kg /day For Systemic Fungal infection - 3-5 mg / kg /day	
2	Anicord	Anidulafungin	Anicord Injection 100 mg	Candidemia and other forms of Candida infections (intra-abdominal abscess and peritonitis), Esophageal candidiasis in adults	Candidemia - 200 mg loading dose on Day 1, followed by 100 mg once daily maintenance dose for at least 14 days after the last positive culture Esophageal candidiasis - 100 mg loading dose on Day 1, followed by 50 mg once daily maintenance dose for a minimum of 14 days and for at least 7 days following resolution of symptoms	
3	Caspocon	Caspofungin Acetate	Caspocon Injection 50 mg Caspocon Injection 70 mg	Empirical therapy for presumed fungal infections in febrile neutropenic patients Treatment of Candidemia and the following Candida infections: Intra-abdominal abscesses, Peritonitis and Pleural space infections Treatment of Esophageal Candidiasis Treatment of invasive Aspergillosis in patients who are refractory to or intolerant of other therapies	70 mg loading dose, followed by 50 mg daily as maintenance dose	
4	Isacrest	Isavuconazole	Isacrest Capsule 100 mg	Treatment of Invasive Aspergillosis and Invasive Mucormycosis (for patients 18 years of age and older) Early targeted therapy (pre-emptive or diagnostic-driven therapy) may be instituted pending confirmation of the disease from specific diagnostic tests.	Loading dose - Two capsules (equivalent to 200 mg of Isavuconazole) every 8 hours for the First 48 hours (6 administrations in total) Maintenance dose - Two capsules (equivalent to 200 mg of Isavuconazole) once daily thereafter	

*Indication & Dose adapted from reference PI

Critical Care

Antifungal Portfolio

CONCORD BIOTECH

Sr. No.	Brand Name	Molecule	SKUs	INDICATION*	DOSE*	PACKSHOT
5	Micacord	Micafungin	Micacord Injection 50 mg Micacord Injection 100 mg	Treatment of patients with Candidemia, Acute Disseminated Candidiasis, Candida Peritonitis and Abscesses Prophylaxis of Candida Infections in Patients Undergoing Hematopoietic Stem Cell Transplantation (HSCT) Treatment of patients with Esophageal Candidiasis	Prophylaxis in HSCT - 50 mg once daily Candidemia, Acute Disseminated Candidiasis, Candida Peritonitis and Abscesses - 100 mg once daily Esophageal Candidiasis - 100 or 150 mg once daily	
6	Picocord GR	Posaconazole	Picocord GR Tablet 100 mg Picocord GR Tablet 300 mg	Prophylaxis of invasive Aspergillus and Candida infections in patients who are at high risk of developing these infections due to being severely immunocompromised	Loading dose – 300mg twice a day on the first day Maintenance dose – 300mg once a day	
7	Vorixia	Voriconazole	Vorixia Tablet 200 mg Vorixia Injection 200 mg	Treatment of Invasive Aspergillosis Treatment of Candidaemia in non-neutropenic patients Treatment of fluconazole-resistant serious invasive Candida infections Prophylaxis of invasive fungal infections in high risk allogeneic hematopoietic stem cell transplant (HSCT) recipients.	Tablet: 200 mg every 12 h >40 kg and 100 mg every 12 h for <40kg Injection: Loading Dose - 6 mg/kg every 12 hours Maintenance Dose - 3-4 mg/kg every 12 hours	

*Indication & Dose adapted from reference PI