Critical Care

Other Products

Anticoagulant Product

| Sr. No. | Brand Name | Molecule | SKUs | INDICATION | DOSE | PACKSHOT |
|------------|------------|------------|--|--|---|--|
| 1 | Anxacord | Enoxaparin | Anxacord PFS 40 mg Anxacord PFS 60 mg | Prophylaxis of venous thromboembolic disease, Treatment of DVT and PE, Treatment of unstable angina and NSTEMI | Prophylaxis of venous thromboembolic disease - 2,000 IU (20 mg) SC once daily Treatment of DVT and PE - 100 IU/kg (1 mg/kg) body weight SC once daily Treatment of unstable angina and NSTEMI - 100 IU/kg (1 mg/kg) body weight SC once daily | Enoxaparin Sodium Injection IP 60 mg/0.6 ml an X-apard -60 grand grand de stocker rayer stated Enoxaparin Sodium Injection IP 40 mg/0.4 ml Enoxaparin Sodium Injection IP 40 mg/0.4 ml Enoxaparin Sodium Injection IP 40 mg/0.4 ml an X-apard -40 grand grand de stocker rayer stated |

Parenteral Nutrition Product

| 2 | Conthion | Glutathione | Conthion Injection 600 mg | Alcoholic Fatty Liver, Alcoholic Liver Fibrosis, Alcoholic Cirrhosis, Alcoholic Hepatitis | In Mild to moderate disease, 300mg to 600mg daily by slow intramuscular or intravenous injection | CONCORD BIOTECT Glutathione for Injection 600 mg Conthion Write sire Lycycliked For IMAILY, use only 1 Correlapas. |
|---|----------|-------------|------------------------------|---|---|---|
|---|----------|-------------|------------------------------|---|---|---|

Critical Care

Other Products

Plasma Products

| Sr. No. | Brand Name | Molecule | SKUs | INDICATION. | DOSE | PACKSHOT |
|------------|-------------------|--------------------------------------|------------------------|---|---|---|
| 3 | Gamacon | Human Normal Immunoglobulin 5% | Gamacon 5gm / 100ml | Primary immunodeficiency, Septic Shock, Kawasaki disease | Primary immunodeficiency - 300-600mg/kg every 3-4 week Septic Shock - 5 gm/day for 3 days or 15g/day for 1 day Kawasaki disease - 400mg/kg daily for 5 consecutive Days | CONCORD BIOTECH CONCORD BIOTECH Human Normal Immunoglobulin IP 5% Gamacon TITHICIT For IV. Usin Markindialtied from Challed frimum Panna Challed frimum Panna Challed frimum Panna Challed frimum Panna So of the Concord |
| 4 | Obulin | Human Normal Albumin 20% | Obulin 20gm / 100ml | Management of Hypovolemia, Hypoalbuminemia | Initial infusion of 100ml, additional amounts may be administered as clinically indicated | CONCORD BIOTECH Aneally Nacional Particular State of the Control |