



Rheumatology

Sr. No.	Brand Name	Molecule	SKUs	INDICATION	DOSE*	PACKSHOT
1	Adacord	Adalimumab	40 mg/0.8 ml PFS	Rheumatoid Arthritis including Juvenile Idiopathic Arthritis, Ankylosing Spondylitis, Psoriatic Arthritis	- Rheumatoid Arthritis, Ankylosing Spondylitis, Psoriatic Arthritis: Every other week Juvenile Idiopathic Arthritis: 15 kg(33 lbs) to <30 kg(66 lbs): 20 mg every other week >30 kg (66lbs): 40 mg every other week	
2	Arthimide	Leflunomide	10 mg Tablets 20 mg Tablets	Rheumatoid Arthritis	Recommended Maintenance dose is 10 mg – 20 mg once daily depending upon the severity of the disease	
3	Tofajoint	Tofacitinib	5 mg Tablets	Rheumatoid Arthritis, Psoriatic Arthritis, Ulcerative Colitis & Polyarticular Course Juvenile Idiopathic Arthritis	Rheumatoid Arthritis: 5 mg twice daily	
4	Tofajoint ER	Tofacitinib Extended Release	11 mg Tablets	Rheumatoid Arthritis, Psoriatic Arthritis, Ulcerative Colitis & Polyarticular Course Juvenile Idiopathic Arthritis	Rheumatoid Arthritis: Once daily dosing	

*Indication & Dose adapted from reference PI

Sr. No.	Brand Name	Molecule	SKUs	INDICATION	DOSE*	PACKSHOT
5	Conimab	Rituximab	500 mg/50 ml Infusion 100 mg/10 ml Infusion	Rheumatoid Arthritis	Administer Rituximab two-1000mg IV infusions separated by 2 weeks	
6	Unuric	Febuxostat	40 mg Tablets 80 mg Tablets	Chronic management of hyperuricemia in patients with gout	Recommended at 40 mg or 80 mg once daily	
7	Coni-Medrol	Methylprednisolone Sodium Succinate Inj.	40 mg/Vial 125 mg/Vial 500 mg/Vial	Rheumatoid Arthritis including Juvenile Rheumatoid Arthritis, Acute Gouty Arthritis, Psoriatic Arthritis, Ankylosing Spondylitis, SLE (& Lupus Nephritis)	Rheumatoid Arthritis: Low-dose glucocorticoids (≤ 10 mg/day of prednisone or equiv. or ≤ 8 mg/day methylprednisolone*) in patients with moderate or high RA disease activity when starting DMARDs & in patients with DMARD failure or biologic failure Disease flares & major extra-articular manifestations may be managed with higher, or even pulse doses (Inj. methylprednisolone 500 mg to 1 g IV daily $\times 3$ doses of glucocorticoids)	

*Indication & Dose adapted from reference PI

Rheumatology

CONCORD BIOTECH

Sr. No.	Brand Name	Molecule	SKUs	INDICATION	DOSE*	PACKSHOT
8	Etoricon	Etoricoxib	• 60 mg Tablets • 90 mg Tablets • 120 mg Tablets	Symptomatic relief of Rheumatoid Arthritis, Ankylosing spondylitis, Osteoarthritis, and the pain and signs of Inflammation associated with acute gouty arthritis	Rheumatoid Arthritis & Ankylosing spondylitis: Recommended Dose is 60 mg once daily Acute Gouty Arthritis: Recommended dose is 120 mg once daily	

*Indication & Dose adapted from reference PI