


Portfolio

Sr. No.	Brand Name	Molecule	SKUs	INDICATION*	DOSE*	PACKSHOT
1	Tacrocord	Tacrolimus 0.1% w/w Tacrolimus 0.03%	Tacrocord Ointment 10 g Tacrocord Ointment 20 g	Moderate-to-severe atopic dermatitis in adults and adolescents (16 years of age and above) Moderate-to-sever atopic dermatitis in children aged 2 to 16 years of age	Flare Treatment: Tacrolimus 0.1% ointment twice a day and treatment should be continued until clearance of the lesion. Maintenance Treatment: Tacrolimus 0.1% ointment to applied once a day, twice weekly. Flare Treatment: Tacrolimus 0.03% ointment twice a day and treatment should be continued until clearance of the lesion. Maintenance Treatment: Tacrolimus 0.03% ointment to applied once a day, twice weekly.	
2	Picroda	Pimecrolimus 1% w/w	Picroda Cream 10 g	Mild-to-moderate atopic dermatitis in patients 3 months of age or older	Apply a thin layer of Pimecrolimus 1% ointment to the affected skin twice daily and rub in gently and completely. Each affected region of the skin should be treated with pimecrolimus until clearance occurs and then treatment should be discontinued. It may be used on all skin areas, including the head and face, neck and intertriginous areas, except on mucous membranes and should not be applied under occlusion.	
3	Jakoway	Tofacitinib 2% w/w	Jakoway Ointment 15 g Jakoway Ointment 30 g	Atopic dermatitis in adults	Apply a thin layer of Tofacitinib 2% ointment twice daily to affected areas.	
4	Hypivent	Desonide 0.05% w/w	Hypivent Cream 30 g	Inflammatory and pruritic manifestations of corticosteroid-responsive dermatoses	Apply to the affected area as a thin film two to four times daily depending on the severity of the condition.	

*Indication & Dose adapted from reference PI