

Dermatology Care

Portfolio

Sr. No.	Brand Name	Molecule	SKUs	INDICATION*	DOSE*	PACKSHOT
1	Muprevent	Mupirocin 2% w/w	Muprevent Ointment 5 g Muprevent Ointment 15 g	Topical treatment of impetigo due to Staphylococcus aureus and Streptococcus pyogenes	Apply a small amount of Mupirocin 2% ointment with cotton swab or gauze pad to the affected area 3 times daily for upto 10 days.	
2	Fusivent	Fusidic acid 2% w/w	Fusivent Cream 5 g Fusivent Cream 15 g	Topical treatment of primary and secondary skin infections	Apply a thin layer of Fusidic Acid 2% cream to the affected area 2-3 times a day, depending on the severity of the infection. Treatment should typically last for 7-10 days, or as directed by the healthcare provider.	
3	Pobix	Neomycin 3400 IU + Polymyxin B Sulfate 5000 IU + Bacitracin Zinc 400 IU	Pobix Ointment USP 20 g	As a first aid to help prevent infections in minor scrapes, cuts and burns	Apply a small amount of this product (an amount equal to the surface area of the tip of a finger) on the area 1 to 3 times daily. Can be covered with a sterile bandage.	

*Indication & Dose adapted from reference PI